

Sheikh Nawaz ALI<sup>1,2\*</sup>, Jyotsna DUBEY<sup>1,3</sup>, Prachita ARORA<sup>1,2</sup>, Shubhra SHARMA<sup>4</sup>,  
P. MORTHEKAI<sup>1,2</sup>, Anupam SHARMA<sup>1,2</sup>, Pankaj KUMAR<sup>5</sup>, Vaibhava SRIVASTAVA<sup>3</sup>

## Glacier sensitivity to climate variability since MIS-2: insights from monsoon dominated north-eastern Himalaya

**Abstract:** Ali S.N., Dubey J., Arora P., Sharma S., Morthekai P., Sharma A., Kumar P., Srivastava V., *Glacier sensitivity to climate variability since MIS-2: Insights from monsoon dominated north-eastern Himalaya*. (IT ISSN 0391-9838, 2024). The present study investigates the glacial landform evolution using moraine stratigraphy in the Lashar and the Chopta valleys (Thangu) of north Sikkim, eastern Indian Himalaya. The older moraines and the outwash sediments have been dated using luminescence and radiocarbon dating techniques which has allowed an understanding of the causes, and patterns of glacier advances/retreat. Chrono-stratigraphy of multiple glacier advances reveals that the oldest glacier advance preserved in the region corresponds to the Marine Isotopic Stage-2 (MIS-2; global Last Glacial Maximum - gLGM) while the subsequent advances are younger. Our data provides a record of four glacier advances, named Thangu Glacier Advances (TGA-I to TGA-IV). The optically stimulated luminescence (OSL) dating has yielded an age range between  $28 \pm 2.8$  and  $22 \pm 2.4$  ka for the oldest glacier advance (TGA-I) dated. The second prominent glacier advance (TGA-II) is correlated with the Younger Dryas cooling event based on pro-glacial sediment OSL ages ranging between  $11 \pm 0.8$  and  $9.9 \pm 0.8$  ka. A minor advance/standstill, TGA-III seems to coincide with the 8.2 ka cooling event. TGA-IV is represented by prominent terminal moraines close to the present day glaciers and would have possibly occurred during the mid-late Holocene cooling. Post-gLGM deglaciation is dated between  $18 \pm 2.2$  and  $15 \pm 2.6$  ka whereas the second deglaciation following TGA-II represented by pro-glacial lake sediments abutting the TGA-II moraines, is dated between  $11 \pm 0.8$  and  $9.9 \pm 0.8$  ka. A close correspondence of glacier advances with periods of temperature minima would imply that the glaciers have actively responded to cooler temperatures associated with the mid-latitude westerlies in response to global variations and retreated during the enhanced monsoonal phases.

**Key words:** Sikkim Himalaya, Last Glacial Maximum, Optical ages, Mid-latitude westerlies, Indian Summer Monsoon.

### INTRODUCTION

Recent studies indicate that the Himalayan glaciers are melting at an alarming rate (Bolch *et al.*, 2012; Lutz *et al.*, 2014; Azam *et al.*, 2018; Maurer *et al.*, 2019), which is likely to affect the hydrology, biodiversity, and ecosystem services in the mountain as well as downstream regions (Buytaert *et al.*, 2017; Milner *et al.*, 2017). These shrinking glaciers provide striking evidence of glacier-climate interactions and how climate shapes the planet Earth (Mote and Kaser, 2007). Therefore, studies pertaining to

modern glacier retreats and comparisons with the past changes in glacier boundary conditions are crucial, as they provide a benchmark for assessing the future response of glaciers to climate change (Solomina *et al.*, 2008). This may be achieved by resolving the timing of glaciation and understanding the role of different weather systems in driving glacier advances and retreats (glacier sensitivity to regional and global climate change). It has been observed that there is a spatial heterogeneity within Himalayan glaciers which is generally attributed to orography and geographic location that modulates the main drivers of glacier mass balance, temperature and precipitation. This heterogeneity is manifested in spatial variability in glacier ice volume and glacier lengths (Benn and Owen, 1998; Sharma and Shukla, 2018). The role of the Indian Summer Monsoon (ISM) and Mid-latitude westerlies (MLWs), is suggested in dictating the pattern of variability of the northwestern Himalayan glaciers (Bookhagen and Burbank, 2010; Murari *et al.*, 2014) how-

<sup>1</sup> Birbal Sahni Institute of Palaeosciences, Lucknow, 226007 India.

<sup>2</sup> Academy of Scientific & Innovative Research (AcSIR), Ghaziabad, 201002 India.

<sup>3</sup> Department of Geology, Institute of Science, Banaras Hindu University, Varanasi, 221005 India.

<sup>4</sup> Physical Research Laboratory, Ahmedabad, 380009 India.

<sup>5</sup> Inter-University Accelerator Centre (IUAC), New Delhi, 110067 India.

\*Corresponding author: Sheikh Nawaz Ali (snawazali@gmail.com)

ever, understanding of the climate dynamics is still at an embryonic stage in the Indian eastern Himalaya (Ali *et al.*, 2019).

Previous studies have suggested that the Himalayan glaciers were out of phase with the global cooling events [e.g., Last Glacial Maximum (LGM)], but instead were sensitive to the variations in the ISM precipitation, particularly in the monsoon-dominated sectors where the response was suggested to be asynchronous (Scherler *et al.*, 2010; Murari *et al.*, 2014). A few, recent studies particularly from the monsoon-dominated western and central Himalaya eloquently demonstrated that the glaciers expanded synchronously with the LGM, and the temperature seems to be an overriding factor in driving the glacier advances (e.g. Ali *et al.*, 2013, 2022; Bali *et al.*, 2013; Eugster *et al.*, 2016; Bisht *et al.*, 2017; Rana *et al.*, 2019). Further, the semi-arid Trans-Himalayan glaciers are suggested to respond more sensitively to moisture availability in comparison to glaciers in relatively humid regions like the central Himalaya which are more sensitive to temperature changes (Zech *et al.*, 2009). However, these suggestions remain inconclusive, as the existing glacial chronological data shows heterogeneity, and hence identifying the climatic controls on glacier fluctuations during the late Quaternary period are still debated (Dortch *et al.*, 2013; Eugster *et al.*, 2016). The present study explores the eastern Himalaya (Sikkim) where moisture is supplied dominantly by the ISM with a subordinate contribution from the MLW. Being located at the leeward side of the orography (higher Himalayas) the archives explored are in the transient climate zone and therefore, should represent the response of the terrain to minor climate fluctuations. Climatologically, the transient climate zone in the Himalayas refers to the region where the influences of the Indian Summer Monsoon (ISM) from the south and mid-latitude westerlies (MLWs) from the north converge. Variations in either of these weather systems significantly impact the local environment, particularly glaciers, which are highly sensitive to even minor changes in weather conditions. Studies suggest that changes in climatic parameters are often amplified in transitional climate zones, making these regions critical for monitoring and understanding climate fluctuations (Fu *et al.*, 1992; Zhou *et al.*, 2016; Ali *et al.*, 2018, 2020). Further, there could be contributions from East Asian summer and winter monsoons which need to be explored further (Gao *et al.*, 2014; Ali *et al.*, 2019, 2020). Therefore, ascertaining the behavior of the glaciers to climate variability and the role of different climate systems is crucial (Herzschuh, 2006; Chen *et al.*, 2008; Maher, 2008). The objectives of the study are (i) to build the morpho-chrono-stratigraphy of the glacier advances/retreat in the region, and (ii) to ascertain the response of the glaciers to climate variability induced by different climate systems in the region.

## REGIONAL SETTINGS

The state of Sikkim is sandwiched between Nepal and Bhutan on the east and west, respectively, and shares its northern border with the Tibetan plateau. Covering an area of  $\sim 7096 \text{ km}^2$ , Sikkim forms the upper catchment of the Tista river basin which is a major tributary of the Brahmaputra River. The state has a steep gradient with an altitudinal range of 280-8568 m asl. The Higher Himalayan sector of Sikkim has 449 glaciers covering an area of  $706 \text{ km}^2$  with an estimated ice volume of  $39.61 \text{ km}^3$  (Raina and Srivastava, 2008). Geographically, the northern district of Sikkim (Mangan) holds the majority of the glaciers and has preserved a suite of landforms documenting past glacier activity.

In the present study three tributary valleys of the upper Tista river catchment (Thangu region), higher (north) Sikkim Himalaya ( $27^{\circ}03'41''$  to  $28^{\circ}7'34''$  N;  $88^{\circ}03'40''$  to  $88^{\circ}57'19''$  E; fig. 1) have been studied due to the availability of evidence of past landscape changes in the form of glacial and glacio-fluvial deposits. Well-preserved moraines, and relict lake sediments (glacio-fluvial) attest to multiple glacier advances in the region. Geologically, the area is located between the South Tibetan Detachment Zone (STDZ) and the Main Central Thrust (MCT) (Dasgupta *et al.*, 2004). The lithology is dominated by the Kanchenjunga/Darjeeling Gneiss (Basu, 2013). The Climatic Research Unit (CRU TS.4.03,  $0.5^{\circ}$  latitude x  $0.5^{\circ}$  longitudes, 1901-2015 CE) reanalysis data suggests that for the grid cell  $27.75$  N  $88.75$  E that encompasses the study area, the mean annual temperature (MAT) of the area is  $\sim 1.5^{\circ}\text{C}$  and the mean annual precipitation (MAP) is  $\sim 700$  mm. Further, the limited *in situ* meteorological data suggest an average precipitation of  $\sim 820$  mm at Thangu (ENVIS, 2016). The study area lies on the southern margin of the Tibetan plateau, which splits the dry westerlies into the northern and southern branches (Tian *et al.*, 2007; Yihui and Zunya, 2008). The latter picks up moisture from the Arabian Sea and continental recycling that can favor significant cooling and increased precipitation, thus promoting snowfall and snow accumulation across the Himalayan-Tibetan orogen (Pang *et al.*, 2012; Gao *et al.*, 2014; Bao and You, 2019). The higher Sikkim Himalaya is dominantly influenced by the ISM ( $>1000$  mm/yr), however, during the winter, the precipitation is facilitated by the eastward transport of water vapor by the southern branch of MLW (Curio *et al.*, 2014; Ali *et al.*, 2019, 2020). Soheb *et al.* (2015) opined that the eastern Himalayan glaciers are summer accumulation type; however, according to Wang *et al.* (2017) the accumulation of the eastern Himalayan glaciers is uncertain and glaciers accumulate mass both in winter and summer, while ablate in late spring. There is a paucity of *in situ* glacier mass balance measurements; however, remote sensing-based estimates (2002-2011) suggest maximum snow accumulation between November and February for which the MLWs are implicated (Krishna, 2005; Basnett and Kulkarni, 2019).




Figure 1 - Map compilation showing (a) the location of the study area i.e. the Sikkim Himalaya, northeast India (yellow box) in the north of the Bay of Bengal (BoB) along with the trajectories of the two major weather systems i.e. the Indian summer monsoon (ISIM) and the mid-latitude westerlies (MLW) with the trajectory of Winter monsoon (WM); (b) Shuttle Radar Topographic Mission (SRTM) digital elevation model (DEM) of Sikkim showing the location of the study area Thangu valley (outlined in yellow) along with the glaciers, major drainages and the two tectonic boundaries i.e. South-Tibetan Detachment System (STDS) in the north and Main Central Thrust (MCT) in the south of the study area.

## MATERIALS AND METHODS

Detailed field surveys supported by the Landsat ETM+ data, Google Earth Pro Imagery, and a three arc-second (~90 m) Shuttle Radar Topography Mission (SRTM) digital elevation model (DEM) have enabled us to construct geomorphological maps. Detailed descriptions are given in our earlier work (Ali *et al.*, 2019; Dubey *et al.*, 2019). Based on the moraine stratigraphy, four moraine building events have been identified in the study area and suggest a widespread glacial cover with large outlet tributary glaciers in the oldest stage. The disposition of the lateral and terminal moraines has been used to estimate the extent of gla-

cial cover during different stages and their corresponding ice volume. We have collected samples for luminescence dating from sand lenses embedded within the moraines as well as from the relict pro-glacial lake sediments abutting the terminal latero-frontal moraine. The sand lenses (ice contact sediments) are present within the moraines and are better bleached as these sediments are generally transported by supraglacial melt-outs and hence receive adequate sunlight and have successfully been utilized for dating moraines (Phillips *et al.*, 2000; Kessler *et al.*, 2012; Ali *et al.*, 2013, 2022). Standard sample preparation procedures for the luminescence dating technique have been followed (Mortheikai and Ali, 2014). The OSL measurements were performed on an automated Risø TL-OSL reader (TL/OSL-DA-20; (Bøtter-Jensen *et al.*, 2010) at the Birbal Sahni Institute of Palaeosciences, Lucknow India. The equivalent doses (De) measurements were carried out on single aliquot regeneration (SAR) protocol (Murray and Wintle, 2000). To avoid any contribution from the feldspar signal (sporadic presence) we applied a post-IR blue during which the aliquots were exposed to IR diodes for 100 s (seconds) at room temperature to suppress the feldspar signal prior to OSL measurements. The elemental concentration measurements for radioactive elements (uranium, thorium, and potassium) for environmental dose rate estimation were carried out using the high-purity Germanium detector (HPGe). Cosmic ray contributions were calculated using the standard method of the contribution of cosmic rays to the dose rates (Prescott and Hutton, 1994). The calculation of OSL ages was done on the Dose Rate and Age Calculator (DRAC) which is an open-access, web-based program (Durcan *et al.*, 2015).

## RESULTS

The study identifies four glacier phases with progressively decreasing magnitude which are named as Thangu glacier Advances (TGA)-I (oldest) to TGA-IV (youngest) (figs 2-3). The oldest and most extensive TGA-I lateral moraines terminated at the exit of the Lashar, Chopta (figs 2a, and Kalip tributary valleys at elevations of ~3850, 4000, and 3700 m asl respectively (Ali *et al.*, 2019; Dubey *et al.*, 2019). The presence of fluvio-glacial lake deposit at the confluence of the tributary and trunk valley suggests that the tributary glaciers extended down to the confluence of the trunk valley and obstructed the meltwater flow – giving rise to glacio-fluvial lake sequences (Dubey *et al.*, 2019; fig. 2a-c). Recessional moraines and glacio-fluvial lake deposits associated with TGA-I manifest the first event of deglaciation (figs 2b-c). The subsequent glacier advances showed a progressive decline in the ice volume and, thus were restricted within the tributary valleys (figs 2-3). The TGA-II moraines have sub-rounded crests and are covered



Figure 2 - Field photographs showing the characteristic lateral/latero-frontal (dashed lines) moraines of Chopta and Lashar valley. (a) left lateral moraine (magenta) representing the Thangu glacier advance-I (TGA-I); (b) close-up of the fluvio-lacustrine sediments formed during TGA-I; (c) recessional moraine representing the first de-glaciation (post-TGA-I) in the region; (d) Right and left lateral moraine of TGA-II in Lashar valley; (e) fluvio-lacustrine sediments formed during TGA-I de-glaciation; (f) moraines of TGA-II, III (downstream view) in the Lashar valley, (g-h) Youngest latero-frontal moraine (TGA-IV) damming the present day pro-glacial lake.

by Juniper shrubs and grasses and are represented by an assemblage of moraine ridges, which are interpreted as multiple standstill phases. These moraines are preserved in the Lashar valley and can be traced down to ~12 km from the contemporary glacier and terminates at ~4300 m asl. The terminal moraine of TGA-II has preserved relict moraine dammed lake sediments (abutting the TGA-II) and represents the second phase of deglaciation (figs 2d-e). The TGA-III is represented by lateral moraine ridge abutting and cutting through the TGA-II lateral moraines and terminates at ~4450 m asl (~10 km from contemporary glacier; fig. 2f). The youngest TGA-IV glacier advance is represented by curvilinear moraine ridges damming the present-day proglacial lakes in the Lashar valley. These moraines are

unstable, comprised of huge boulders, sharp-crested, and devoid of any vegetation (figs 2g-h).

Twelve ages, which include 8 published (Ali *et al.*, 2019) and 3 new OSL ages and 1 radiocarbon age (Ali *et al.*, 2018) form the chrono-stratigraphic framework for glacier advances and retreats (fig. 3; table 1). The oldest TGA-I is dated to  $28 \pm 2.8$  and  $22 \pm 2.4$  ka. An age of  $6 \pm 1.7$  ka has been removed being an outlier. The TGA-II moraines are associated with relict proglacial lake sediments, which have been dated to  $11 \pm 0.8$ ,  $10.7 \pm 1.2$ , and  $9.9 \pm 0.8$  ka, implying deglaciation during ameliorating climatic conditions. The TGA-III and IV moraines remain undated due to the scarcity of the datable material which is sand lenses embedded within the moraine.




Table 1 - A table providing the details of luminescence ages used in the present study.

Glacier phase	Sample code	De (Gy)	U (ppm)	Th (ppm)	K %	Dose rate (Gy/ka)	Age (ka)
TGA - I (glacier advance)	CH OSL-5	99 ± 3	2.3	12.2	2.2	3.6 ± 0.3	28 ± 2.8
	CH OSL -6	97 ± 3	3	11.2	2.7	3.9 ± 0.3	25 ± 2.6
	CH OSL -7	24 ± 2	4.1	14.6	2.99	4.0 ± 0.3	6 ± 2.4
	CH OSL -8	70 ± 3	2.8	13	2.6	3.2 ± 0.2	22 ± 2.4
TGA - I (de-glaciation)	CH OSL -1	74 ± 4	5.6	23.6	2.6	4.8 ± 0.3	15 ± 2.6
	CH OSL -2	72 ± 4	4.2	16.8	2.6	4.2 ± 0.3	17 ± 2.8
	CH OSL -3	65 ± 2	3.8	18.1	2.6	4.0 ± 0.3	16 ± 2.6
	CH OSL -4	59 ± 2	2.7	12.5	2.5	3.3 ± 0.2	18 ± 2.2
TGA - II (glacier advance)	Younger Dryas						
TGS - II (de-glaciation)	TV OSL -1	50 ± 5	4.7	14.8	2.6	4.5 ± 0.2	11.0 ± 0.8
	TV OSL -2	45 ± 2	5	14.1	2.2	4.2 ± 0.2	10.7 ± 1.2
	TV OSL -3	51 ± 3	5.1	22.6	2.8	5.2 ± 2.6	9.9 ± 0.8
TGA - III (glacier advance)	Early mid Holocene (?)						
TGA - IV (glacier advance)	Mid-late Holocene (?)						

## DISCUSSION

The dynamics of the late Quaternary glaciation in the Himalaya, especially in the ISM-dominated regions, advocate for a complex interplay of insolation-induced changes in the ISM and MLW (Benn and Owen, 1998; Finkel *et al.*, 2003; Sati *et al.*, 2014). Various studies from the Himalayan region suggest either asynchronous glacier response to global climate forcings (Benn and Owen, 1998; Rupper *et al.*, 2009; Scherler *et al.*, 2010) or in sync response of the glaciers with the Northern Hemisphere ice-sheet dynamics and global climate perturbations (Ali *et al.*, 2013; Sati *et al.*, 2014; Bisht *et al.*, 2017; Rana *et al.*, 2019). The Upper Tista valley is dominantly influenced by the Bay of Bengal branch of the ISM (annual precipitation  $>1000$  mm); however, its specific location facilitates the penetration of the mid-latitude westerlies characterized by winter precipitation that occurs in the form of snowfall during December to March (Ali *et al.*, 2019) which therefore makes the region critical to understand the boundary conditions of ISM and MLW in driving the glacier response.

Integrating the geomorphological and chronological results, it is evident that during the early MIS-2 including the gLGM (TGA-I;  $28 \pm 2.8$  and  $22 \pm 2.4$  ka), the upper Tista valley was occupied by large tributary glaciers that extended up to 16 km1, which is similar in magnitude to other ISM influenced Himalayan sectors (Ali *et al.*, 2013, 2019, 2022; Bisht *et al.*, 2017; Shukla *et al.*, 2018; Peng *et al.*, 2019; Rana *et al.*, 2019; Kumar *et al.*, 2023) (fig. 4). This glacier advance initiated during weaker ISM phase at  $28 \pm 2.8$  which broadly corresponds to Heinrich event 3, when the Laurentide ice sheet was discharging exceptionally large icebergs into north Atlantic resulting in cooler sea surface temperature (Heinrich *et al.*, 1988; Bond *et al.*, 1992, 1993; Hodell *et al.*, 2017; Liu *et al.*, 2020). This would imply that the onset of glaciation was controlled by temperature decline in the Himalayan region (Ganju *et al.*, 2018). This event persisted/sustained till the gLGM under the influence of decreasing solar insolation and ice-albedo feedback mechanism (Adams *et al.*, 1999; Colin *et al.*, 1998). Considering that the ISM was weak during the LGM, it is reasonable to assign the TGA-I to decreased temperatures associated with the enhanced mid-latitude westerlies (Ali *et al.*, 2022; Ganopolski *et al.*, 1998) characterized by increased winter precipitation (Ghosh *et al.*, 2015) and hence snow accumulation on the Himalayan-Tibetan orogen, which is also supported by modern meteorological and meltwater isotopic data (Sirocko *et al.*, 1996; An *et al.*, 2012; Yang and Ding, 2014; DiNezio *et al.*, 2018; Ali *et al.*, 2020). It has been suggested that the temperature in the eastern Himalaya was almost  $4.6^{\circ}\text{C}$  lower than present during the LGM (Yang *et al.*, 2022). This colder climate, that prompted glacier advance,

has been linked to low solar insolation precession cycles and a stronger negative forcing driven mostly by lower levels of atmospheric carbon dioxide (Otto-Thompson *et al.*, 1997; Bliesner *et al.*, 2006; Braconnot *et al.*, 2007). It is also observed that the glacier expansion in eastern Sikkim Himalaya was not as extensive as in westerly dominated Nubra valley ( $\sim 80$  km towards south; Nagar *et al.*, 2013; Ganju *et al.*, 2018) or the Chandra valley ( $\sim 200$  km towards west till Udaipur township; Eugster *et al.*, 2016). Yet, the advancement is comparable to the ISM dominated central Himalaya and climatologically transitional southern Zanskar Himalaya (Ali *et al.*, 2013, 2022; Bisht *et al.*, 2017; Sharma and Shukla, 2018; Shukla *et al.*, 2018; Rana *et al.*, 2019). Although the explanation of a generally restricted but geographically diverse LGM glacier expansion remains unresolved to date, a lower insolation-driven weaker ISM might be the primary reason for the Himalaya's limited LGM glacier expansion (Herzschnuh, 2006; Agrawal *et al.*, 2012; Dutt *et al.*, 2015), while glacier valley orientations, hypsometry, and orography influence may have had a part in non-uniform glacier advances (Yan *et al.*, 2018; Ali *et al.*, 2022).





Figure 4 - Correlation of the present study with other well-established glacial chronologies from similar ISM-dominated Central Himalayan glacier valleys. (a) Present study (Thangu valley, eastern Himalaya; blue-glacier advance, red-glacier retreat); (b) Tons valley (Scherler *et al.*, 2010); (c) Dokriani Glacier valley (Shukla *et al.*, 2018); (d) Kanchenjunga Himalaya (Tsukamoto *et al.*, 2002); (e) Central Himalaya (Murari *et al.*, 2014); (f) Dunagiri valley (Sati *et al.*, 2014); (g) Bhagirathi valley (Barnard *et al.*, 2004a), (h) Goriganga valley (Barnard *et al.*, 2004b).

The recessional moraines representing glacier stabilization during recession/deglaciation are dated between  $18 \pm 2.2$  ka and  $15 \pm 2.6$  ka coinciding with a period of gradually increasing, but weaker ISM (Dutt *et al.*, 2015), and an enhanced westerlies period as inferred from the loess deposit on Tibetan plateau (Li *et al.*, 2016). Within the age uncertainty, the recessional moraine correlates well with the Heinrich (H1) event – which was triggered by large freshwater discharges from the North American ice sheet into the North Atlantic, causing long-lived cold states and implicated for monsoon weakening through the weakening of thermohaline circulation (Heinrich, 1988; Bond *et al.*, 1992; Tierney *et al.*, 2016; Lauterbach *et al.*, 2020). Hence, we propose that the standstill condition during the TGA-I deglaciation was a result of lowered temperatures. Our inferences are supported by the cooler sea surface temperature, and enriched  $\delta^{18}\text{O}_{\text{sw}}$  values (18.2 and 15.6 ka), suggesting weak ISM and lesser outflow of rivers into the Bay of Bengal (Rashid *et al.*, 2011; Liu *et al.*, 2020) (fig. 5). The post gLGM increase in insolation/temperature resulted in a gradual strengthening in the ISM between

15 and 13 ka (Rashid *et al.*, 2011; Dutt *et al.*, 2015; Ghosh *et al.*, 2015; Ali *et al.*, 2018), however the moisture could not transform into solid precipitation (snow) (Rana *et al.*, 2019), thus failed to assist the expansion of valley glaciers. This is also evident from the proglacial peat/bog sequence dated to  $\sim 13 \pm 0.5$  ka suggesting ameliorating climatic conditions in the study area (Ali *et al.*, 2018).

The evidence of TGA-II is present in the tributary Lashar valley (fig. 3). For the TGA-II moraines, a direct age control could not be established due to the paucity of datable (sand lenses) material. However, the ages obtained on the relict pro-glacial lake sediments abutting the TGA-II terminal moraine suggest that the advance is broadly concomitant with the timing of sudden cooling associated with the Younger Dryas event (YD) (12.7-117 ka; Dansgaard *et al.*, 1989; Alley, 2000; Wang *et al.*, 2001) (figs 4-5). This regional cooling is well represented by the more negative values from both NGRIP and Guliya  $\delta^{18}\text{O}$  records (Johnsen *et al.*, 1997; Thompson *et al.*, 1997; Andersen *et al.*, 2006) and associated glacier advances in Himalaya (Mehta *et al.*, 2012; Murari *et al.*, 2014; Hu *et al.*,

Figure 5 - Figure showing correlations between the present data and various palaeoclimatic proxies plotted as a function of age (ka): (a) extent of each glacier advance in terms of distance and position of various glacial landforms in the study area with respect to the present day glacial snout; (b) representation of the areal extent of each glacial advance along with the time frame in accordance with the chronology obtained (unobtained); (c) rank plot showing the age distribution of different Thangu valley glacier advances (blue) and retreat (red), (e)  $\delta^{18}\text{O}_{\text{sw}}$  reconstructed from core BoB-24 (Liu *et al.*, 2020); (d) mean effective moisture variations from the monsoonal Central Asia (Herszschuh, 2006); (e)  $\delta^{18}\text{O}_{\text{sw}}$  reconstructed from core BoB-24 (Liu *et al.*, 2020); (f) speleothem  $\delta^{18}\text{O}$  records from Dongge Cave in the southern China (green; Yuan *et al.*, 2004) and Hulu Cave in the eastern China (red; Wang *et al.*, 2001); (g) Summer insolation data (June insolation 30°N) (Berger and Loutre, 1991).



2015; Ali *et al.*, 2022). During the YD, the eastern Himalaya has witnessed weak ISM conditions with cooler temperatures (Ali *et al.*, 2018) which is also inferred from the sea surface temperature and  $\delta^{18}\text{O}_{\text{sw}}$  values of Bay of Bengal (Kudrass *et al.*, 2001; Liu *et al.*, 2020), enriched  $\delta^{18}\text{O}$  values from Hulu cave stalagmites (Wang *et al.*, 2001) and the Dali Lake (Fan *et al.*, 2019). Glacier expansion during YD (TGA-II) in the eastern Himalaya further highlights the sensitivity of monsoon-dominated glaciers towards millennial-scale cold events. Our suggestions are further corroborated by the equilibrium line altitude (ELA) based temperature reconstruction in the adjacent Zemu and east Rathong valleys wherein a drop of  $\sim 1.7$  °C has been estimated during YD (Kumar *et al.*, 2023). Our observation is at variance with earlier suggestions of insignificant temperature control on glacier response (advance) (Scherler *et al.*, 2010) to rapid climate changes that originated from the North Atlantic and perturbed into the Himalaya (Clark *et al.*, 2002; Denton *et al.*, 2005).

The TGA-II deglaciation ( $11 \pm 0.8$  and  $9.9 \pm 0.8$  ka) is represented by the relict proglacial lake sediments that abut the TGA-II terminal moraine in Lashar valley. Therefore, the early Holocene deglaciation is coeval with the enhanced ISM phase suggested to have spatial influence extending till  $\sim 86^\circ\text{E}$  latitude (Hudson and Quade, 2013). Our inferences get further confidence from the depleted  $\delta^{13}\text{C}$  values of proglacial lake sediments in the study area suggesting enhanced ISM (Ali *et al.*, 2018). These inferences corroborate well with the oxygen isotopic data from the Bay of Bengal and Andaman Sea (Rashid *et al.*, 2011; Liu *et al.*, 2020). Furthermore, the depleted  $\delta^{18}\text{O}$  in speleothem records from northeast India (Berkelhammer *et al.*, 2012; Dutt *et al.*, 2015) and the Tianmen cave (Cai *et al.*, 2012) also indicate insolation driven early Holocene strengthened ISM which began to decline during early to mid-Holocene (Herzschuh, 2006; Cai *et al.*, 2012; Liu *et al.*, 2020). The moraine formation of TGA-II is attributed to the enhanced influence of mid-latitude westerlies (MLWs) during the Younger Dryas (YD) cooling event, while the subsequent deglaciation is associated with abrupt warming, as reported by Ali *et al.* (2018) and further supported by Liu *et al.* (2020) and Rashid *et al.* (2011). The TGA-III remains undated, however based on regional correlation (fig. 5) it may be linked to the early mid-Holocene and a manifestation of the cooling associated with the two closely spaced discharges at  $\sim 8.22$  and  $\sim 8.16$  cal ka BP from glacial lakes Agassiz and Ojibway during the terminal demise of the Laurentide ice sheet (Barber *et al.*, 1999; Rohling and Palike, 2005; Brouard *et al.*, 2021). This event is implicated for the weakening of Atlantic meridional overturning circulation (AMOC), triggering of abrupt 8.2 ka cooling event, and weakening of the ISM (Sinha *et al.*, 2005; Li *et al.*, 2012; Dixit *et al.*, 2014; Rawat *et al.*, 2015). The glacier expansion in response to 8.2 ka cooling, displays the sensitivity of the glaciers to short-lived (centennial-scale) temperature changes (Hahn and Shukla, 1976; Barnett *et al.*, 1989; Colin *et al.*, 1998; Adams *et al.*, 1999; Sharma and Shukla, 2018). We hypothesize that cooler temperatures would have resulted in more winter precipitation, longer winters, and extended cooling, resulting in a positive glacial mass balance and supporting the glacier expansion (Bush, 2002; Ali and Juyal, 2013; Ali *et al.*, 2013, 2019; Murari *et al.*, 2014). The inference would remain tentative until the moraines are dated and coincide with the 8.2 ka cold episode in this region (Scherler *et al.*, 2010; Murari *et al.*, 2014 and references therein).

TGA-IV also remains undated due to the unavailability of datable material for luminescence dating. However, based on similar marginal glacier expansions documented in the ISM-dominated Himalaya (Scherler *et al.*, 2010; BGS-II, Dunagiri valley from Sati *et al.*, 2014, we propose that this glacier advance corresponds to the mid-late Holocene (fig. 5). Climatically, the advance occurred during a weak ISM (Ali *et al.*, 2018), but moderately enhanced westerlies phase characterised by an overall cooling (Wang *et al.*, 2005; Herzschuh, 2006; Otto-Bliesner *et al.*, 2006; Jia *et al.*, 2018) and is correlatable with worldwide glacial advance that occurred at 5.8-4.9 ka (Denton and Karlén, 1973; Shi *et al.*, 1994). These correlations show the greater influence of temperature in driving the glaciers in this region, which itself is facilitated by the enhanced MLW (Herzschuh, 2006).

Contrary to the earlier suggestion of ISM as the major driver of glaciation in the Himalaya (Arora *et al.*, 2023; Finkel *et al.*, 2003; Gayer *et al.*, 2006), recent studies have suggested that the glacier advances since the gLGM were influenced by both precipitation changes and lower temperatures (Rupper *et al.*, 2009; Sati *et al.*, 2014). The combined chronological and field data from north Sikkim suggest that the glaciers advances broadly correspond to negative  $\delta^{18}\text{O}$  values in the Guliya and NGRIP ice core data. Considering that the glaciers showed an advance during weaker ISM phases, we hypothesize that the decline in ISM would have taken place at the expanse of enhanced MLW associated with cooler temperatures and winter precipitation (Wei and Gasse, 1999; Yihui and Zunya, 2008). These phases also correspond to a decline in the solar irradiance outputs (Berger and Loutre, 1991; Steinhilber *et al.*, 2009). Our interpretation gets further confidence from the observation of an overall higher westerly influence during these phases in central Asia (Herzschuh, 2006). With reference to the present data from the eastern Himalaya, the glacier advances appear to be coeval with northern hemisphere cooling which is propagated to the Himalayan region by enhanced MLW reflecting hemisphere-scale processes. Contrary to earlier propositions, the deglaciation is closely linked to increased insolation-driven ISM episodes.

## CONCLUSIONS

The Upper Tista valley is mainly influenced by the ISM; nevertheless, its geographical location allows for the penetration of mid-latitude westerlies, which are characterized by winter precipitation in the form of snowfall. Evidence of four prominent glacier advances (oldest to youngest; TGA-I to IV) extending back to ~28 ka have been reported.

We propose that these glacier advances in the ISM-dominated eastern Himalaya are a manifestation of both precipitation (snowfall) and cooler thermal regime associated with the enhanced MLW, implying that the glaciers in high precipitation areas respond more sensitively to temperature changes.

The increased solid precipitation and propagation of cooler conditions are linked to the weakening of Atlantic meridional overturning circulation (sudden freshwater release from the Laurentide Ice Sheet into the North Atlantic), responsible for an overall cooling in the northern Hemisphere.

Keeping all the above facts in view, it is evident that the glacier advances in the eastern Himalaya are dominantly controlled by MLWs while the subsequent deglaciations are governed by warmer phases (increased insolation).

## ACKNOWLEDGMENTS

The authors are thankful to the Director, BSIP, Lucknow, and Govt. of Sikkim, India. Financial support from DST (SR/DGH-89/2014). We also express our heartfelt thanks to the anonymous reviewers for their insightful comments and constructive feedback, which greatly contributed to improving the quality of this work. Their time and expertise are deeply appreciated.

## AUTHOR CONTRIBUTIONS

S.N. Ali and A. Sharma conceived the study developed the overall methodology and carried out the fieldwork along with J. Dubey. P. Arora and P. Morthekai helped in data analysis. S.N. Ali, S. Sharma, J. Dubey, and P. Arora carried out the writing of the manuscript with the help of all the co-authors. All authors have read and agreed to the published version of the manuscript.

## DATA AVAILABILITY

Data needed will be made available on request.

## REFERENCES

Adams J., Maslin M., Thomas E., 1999. *Sudden climate transitions during the Quaternary*. Progress in Physical Geography, 23 (1), 1-36. <https://doi.org/10.1177/030913339902300101>

Agrawal S., Sanyal P., Sarkar A., Jaiswal M.K., Dutta K., 2012. *Variability of Indian monsoonal rainfall over the past 100 ka and its implication for C3-C4 vegetational change*. Quaternary Research, 77 (1), 159-170. <https://doi.org/10.1016/j.yqres.2011.09.003>

Ali S.N., Biswas R.H., Shukla A.D., Juyal N., 2013. *Chronology and climatic implications of Late Quaternary glaciations in the Goriganga valley, central Himalaya, India*. Quaternary Science Reviews, 73, 59-76. <https://doi.org/10.1016/j.quascirev.2013.05.016>

Ali S.N., Dubey J., Ghosh R., Quamar M.F., Sharma A., Morthekai P., Agrawal S., 2018. *High frequency abrupt shifts in the Indian summer monsoon since Younger Dryas in the Himalaya*. Scientific Reports, 8 (1), 9287. <https://doi.org/10.1038/s41598-018-27597-6>

Ali S.N., Dubey J., Morthekai P., Sharma A., Singh R., Prizomwala S., 2019. *Climate forcing and the initiation of glacier advance during MIS 2 in the North Sikkim Himalaya, India*. Journal of Asian Earth Sciences, 174, 381-388. <https://doi.org/10.1016/j.jseaes.2019.02.005>

Ali S.N., Sharma A., Agrawal S., Yadava M.G., Jani R.A., Dubey J., Morthekai P., 2020. *Oxygen and deuterium isotope characteristics of Teesta river catchment from Sikkim Himalaya, India: implications of different moisture sources*. Geochemical Journal, 54 (5), 327-336. <https://doi.org/10.2343/geochemj.2.0604>

Ali S.N., Singh P., Arora P., Bisht P., Morthekai P., 2022. *Luminescence dating of late pleistocene glacial and glacio-fluvial sediments in the Central Himalaya, India*. Quaternary Science Reviews, 284, 107464. <https://doi.org/10.1016/j.quascirev.2022.107464>

Ali S.N., Juyal N., 2013. *Chronology of late quaternary glaciations in Indian Himalaya: a critical review*. Journal of the Geological Society of India, 82 (6), 628-638. <https://doi.org/10.1007/s12594-013-0201-9>

Alley R.B., 2000. *The Younger Dryas cold interval as viewed from central Greenland*. Quaternary Science Reviews, 19 (1-5), 213-226. [https://doi.org/10.1016/S0277-3791\(99\)00062-1](https://doi.org/10.1016/S0277-3791(99)00062-1)

An Z., Colman S.M., Zhou W., Li X., Brown E.T., Jull A.J., Cai Y., Huang Y., Lu X., Chang H., Song Y., 2012. *Interplay between the Westerlies and Asian monsoon recorded in Lake Qinghai since 32 ka*. Scientific Reports, 2 (1), 1-7. <https://doi.org/10.1038/srep00619>

Andersen K.K., Svensson A., Johnsen S.J., Rasmussen S.O., Bigler M., Röthlisberger R., Ruth U., Sigaard-Andersen M.L., Steffensen J.P., Dahl-Jensen D., Vinther B.M., 2006. *The Greenland ice core chronology 2005, 15-42 ka. Part 1: constructing the time scale*. Quaternary Science Reviews, 25 (23-24), 3246-3257. <https://doi.org/10.1016/j.quascirev.2006.08.002>

Arora P., Ali S.N., Sharma A., 2023. *Role of Different Moisture Sources in Driving the Western Himalayan Past-glacier Advances*. Journal of Atmospheric Science Research, 6 (3), 1-19. <http://doi.org/10.30564/jasr.v6i3.5581>

Azam M.F., Wagnon P., Berthier E., Vincent C., Fujita K., Kargel J.S., 2018. *Review of the status and mass changes of Himalayan-Karakoram glaciers*. Journal of Glaciology, 64 (243), 61-74. <https://doi.org/10.1017/jog.2017.86>

Bali R., Ali S.N., Agarwal K.K., Rastogi S.K., Krishna K., Srivastava P., 2013. *Chronology of late Quaternary glaciation in the Pindar valley, Alaknanda basin, Central Himalaya (India)*. Journal of Asian Earth Sciences, 66, 224-233. <https://doi.org/10.1016/j.jseaes.2013.01.011>

Bao Y., You Q., 2019. *How do westerly jet streams regulate the winter snow depth over the Tibetan Plateau?*. Climate Dynamics, 53 (1), 353-370. <https://doi.org/10.1007/s00382-018-4589-1>

Barber D.C., Dyke A., Hillaire-Marcel C., Jennings A.E., Andrews J.T., Kerwin M.W., Bilodeau G., McNeely R., Southon J., Morehead M.D., Gagnon J.M., 1999. *Forcing of the cold event of 8,200 years ago by catastrophic drainage of Laurentide lakes*. Nature, 400 (6742), 344-348. <https://doi.org/10.1038/22504>

Barnard P.L., Owen L.A., Finkel R.C., 2004a. *Style and timing of glacial and paraglacial sedimentation in a monsoon-influenced high Himalayan environment, the upper Bhagirathi Valley, Garhwal Himalaya*.

Sedimentary Geology, 165 (3-4), 199-221. <https://doi.org/10.1016/j.sedgeo.2003.11.009>

Barnard P.L., Owen L.A., Sharma M.C., Finkel R.C. 2004b. *Late quaternary (Holocene) landscape evolution of a monsoon-influenced high Himalayan valley, Gori-Ganga, Nanda Devi, NE Garhwal*. Geomorphology, 61 (1-2), 91-110. <https://doi.org/10.1016/j.geomorph.2003.12.002>

Barnett T.P., Dümenil L., Schlese U., Roeckner E., Latif M. 1989. *The effect of Eurasian snow cover on regional and global climate variations*. Journal of the Atmospheric Sciences, 46 (5), 661-686.

Basnett S., Kulkarni A.V., 2019. *Snow cover changes observed over Sikkim Himalaya*. In: Saikia A., Thapa P. (Eds), Environmental Change in the Himalayan Region: Twelve Case Studies, 219-232. Springer, Cham. <https://doi.org/10.1007/978-3-030-03362-0>

Basu S.K., 2013. *Geology of Sikkim state and Darjeeling district of West Bengal*. Bangalore, Geological Society of India.

Benn D.I., Owen L.A., 1998. *The role of the Indian summer monsoon and the mid-latitude westerlies in Himalayan glaciation: review and speculative discussion*. Journal of the Geological Society, 155 (2), 353-363. <https://doi.org/10.1144/gsjgs.155.2.0353>

Berger A., Loutre M.F., 1991. *Insolation values for the climate of the last 10 million years*. Quaternary Science Reviews, 10 (4), 297-317. [https://doi.org/10.1016/0277-3791\(91\)90033-Q](https://doi.org/10.1016/0277-3791(91)90033-Q)

Berkelhammer M., Sinha A., Stott L., Cheng H., Pausata F.S., Yoshimura K., 2012. *An abrupt shift in the Indian monsoon 4000 years ago*. In: Giosan L., Fuller D.Q., Nicoll L., Flad R.K., Clift, P.D. (Eds), Climates, Landscapes, and Civilizations, Geophysical Monograph Series, 198 (7), 75-88. <https://doi.org/10.1029/2012GM001207>

Bisht P., Ali S.N., Rana N., Singh S., Sundriyal Y.P., Bagri D.S., Juyal N. 2017. *Pattern of Holocene glaciation in the monsoon-dominated Kosa Valley, central Himalaya, Uttarakhand, India*. Geomorphology, 284, 130-141. <https://doi.org/10.1016/j.geomorph.2016.11.023>

Bolch T., Kulkarni A., Kääb A., Huggel C., Paul F., Cogley J.G., Frey H., Kargel J.S., Fujita K., Scheel M. and Bajracharya S., 2012. *The state and fate of Himalayan glaciers*. Science, 336 (6079), 310-314. <https://doi.org/10.1126/science.1215828>

Bond G., Broecker W., Johnsen S., McManus J., Labeyrie L., Jouzel J., Bonani G., 1993. *Correlations between climate records from North Atlantic sediments and Greenland ice*. Nature, 365 (6442), 143-147. <https://doi.org/10.1038/365143a0>

Bond G., Heinrich H., Broecker W., Labeyrie L., McManus J., Andrews J., Huon, S., Jantschik R., Clasen S., Simet C., Tedesco K., 1992. *Evidence for massive discharges of icebergs into the North Atlantic Ocean during the last glacial period*. Nature, 360 (6401), 245-249. <https://doi.org/10.1038/360245a0>

Bookhagen B., Burbank D.W., 2010. *Toward a complete Himalayan hydrological budget: Spatiotemporal distribution of snowmelt and rainfall and their impact on river discharge*. Journal of Geophysical Research: Earth Surface, 115 (F3). <https://doi.org/10.1029/2009JF001426>

Bötter-Jensen L., Thomsen K.J., Jain M., 2010. *Review of optically stimulated luminescence (OSL) instrumental developments for retrospective dosimetry*. Radiation Measurements, 45 (3-6), 253-257. <https://doi.org/10.1016/j.radmeas.2009.11.030>

Braconnot P., Otto-Bliesner B., Harrison S., Joussaume S., Peterchmitt J.Y., Abe-Ouchi A., Crucifix M., Driesschaert E., Fichefet T., Hewitt C.D., Kageyama M., 2007. *Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum—Part 1: experiments and large-scale features*. Climate of the Past, 3 (2), 261-277. <https://doi.org/10.5194/cp-3-261-2007>

Brouard E., Roy M., Godbout P.M. Veillette J.J., 2021. *A framework for the timing of the final meltwater outbursts from glacial Lake Agassiz-Ojibway*. Quaternary Science Reviews, 274, p.107269. <https://doi.org/10.1016/j.quascirev.2021.107269>

Bush A.B., 2002. *A comparison of simulated monsoon circulations and snow accumulation in Asia during the mid-Holocene and at the Last Glacial Maximum*. Global and Planetary Change, 32 (4), 331-347. [https://doi.org/10.1016/S0921-8181\(02\)00074-7](https://doi.org/10.1016/S0921-8181(02)00074-7)

Buytaert W., Moulds S., Acosta L., De Bievre B., Olmos C., Villacis M., Tovar C., Verbist K.M., 2017. *Glacial melt content of water use in the tropical Andes*. Environmental Research Letters, 12 (11), 114014. <https://doi.org/10.1088/1748-9326/aa926c>

Cai Y., Zhang H., Cheng H., An Z., Edwards R.L., Wang X., Tan L., Liang F., Wang J., Kelly M., 2012. *The Holocene Indian monsoon variability over the southern Tibetan Plateau and its teleconnections*. Earth and Planetary Science Letters, 335, 135-144. <https://doi.org/10.1016/j.epsl.2012.04.035>

Chen F., Yu Z., Yang M., Ito E., Wang S., Madsen D.B., Huang X., Zhao Y., Sato T., Birks H.J.B., Boomer, I., 2008. *Holocene moisture evolution in arid central Asia and its out-of-phase relationship with Asian monsoon history*. Quaternary Science Reviews, 27 (3-4), 351-364. <https://doi.org/10.1016/j.quascirev.2007.10.017>

Clark P.U., Pisias N.G., Stocker T.F., Weaver A.J., 2002. *The role of the thermohaline circulation in abrupt climate change*. Nature, 415 (6874), 863-869. <https://doi.org/10.1038/415863a>

Colin C., Kissel C., Blamart D., Turpin L., 1998. *Magnetic properties of sediments in the Bay of Bengal and the Andaman Sea: impact of rapid North Atlantic Ocean climatic events on the strength of the Indian monsoon*. Earth and Planetary Science Letters, 160 (3-4), 623-635. [https://doi.org/10.1016/S0012-821X\(98\)00116-2](https://doi.org/10.1016/S0012-821X(98)00116-2)

Curio J., Maussion F., Scherer D., 2015. *A 12-year high-resolution climatology of atmospheric water transport over the Tibetan Plateau*. Earth System Dynamics, 6 (1), 109-124. <https://doi.org/10.5194/esd-6-109-2015>

Dansgaard W., White J.W., Johnsen S.J., 1989. *The abrupt termination of the Younger Dryas climate event*. Nature, 339 (6225), 532-534. <https://doi.org/10.1038/339532a0>

Dasgupta S., Ganguly J., Neogi S., 2004. *Inverted metamorphic sequence in the Sikkim Himalayas: crystallization history, P-T gradient and implications*. Journal of Metamorphic Geology, 22 (5), 395-412. <https://doi.org/10.1111/j.1525-1314.2004.00522.x>

Denton G.H., Karlén W., 1973. *Holocene climatic variations, their pattern and possible cause*. Quaternary Research, 3 (2), 155-205. [https://doi.org/10.1016/0033-5894\(73\)90040-9](https://doi.org/10.1016/0033-5894(73)90040-9)

Denton G.H., Alley R.B., Comer G.C., Broecker W.S., 2005. *The role of seasonality in abrupt climate change*. Quaternary Science Reviews, 24 (10-11), 1159-1182. <https://doi.org/10.1016/j.quascirev.2004.12.002>

DiNezio P.N., Tierney J.E., Otto-Bliesner B.L., Timmermann A., Bhattacharya T., Rosenbloom N., Brady E., 2018. *Glacial changes in tropical climate amplified by the Indian Ocean*. Science Advances, 4 (12), eaat9658. <https://doi.org/10.1126/sciadv.aat9658>

Dixit Y., Hodell D.A., Sinha R., Petrie C.A., 2014. *Abrupt weakening of the Indian summer monsoon at 8.2 kyr BP*. Earth and Planetary Science Letters, 391, 16-23. <https://doi.org/10.1016/j.epsl.2014.01.026>

Dortch J.M., Owen L.A., Caffee M.W., 2013. *Timing and climatic drivers for glaciation across semi-arid western Himalayan-Tibetan orogen*. Quaternary Science Reviews, 78, 188-208. <https://doi.org/10.1016/j.quascirev.2013.07.025>

Dubey J., Ali S.N., Sharma A., Morthekai P., Singh R., Sharma R.K., Pandey P., Thakur B., Srivastava V., 2019. *Glacial geomorphology and landscape evolution of the Thangu valley, North Sikkim Himalaya, India*. Journal of the Indian Society of Remote Sensing, 47 (5), 821-837. <https://doi.org/10.1007/s12524-019-00943-5>

Durcan J.A., King G.E., Duller G.A., 2015. *DRAC: Dose Rate and Age Calculator for trapped charge dating*. Quaternary Geochronology, 28, 54-61. <https://doi.org/10.1016/j.quageo.2015.03.012>

Dutt S., Gupta A.K., Clemens S.C., Cheng H., Singh R.K., Kathayat G., Edwards R.L., 2015. *Abrupt changes in Indian summer monsoon strength during 33,800 to 5500 years BP*. Geophysical Research Letters, 42 (13), 5526-5532. <https://doi.org/10.1002/2015GL064015>

Eugster P., Scherler D., Thiede R.C., Codilean A.T., Strecker M.R., 2016. *Rapid Last Glacial Maximum deglaciation in the Indian Himalaya coeval with midlatitude glaciers: new insights from 10Be-dating of ice-polished bedrock surfaces in the Chandra Valley, NW Himalaya*. Geophysical Research Letters, 43 (4), 1589-1597. <https://doi.org/10.1002/2015GL066077>

Fan J., Xiao J., Wen R., Zhang S., Huang Y., Yue J., Wang X., Cui L., Li H., Xue D., Liu Y., 2019. *Mineralogy and carbonate geochemistry of the Dali Lake sediments: Implications for paleohydrological changes in the East Asian summer monsoon margin during the Holocene*. Quaternary International, 527, 103-112. <https://doi.org/10.1016/j.quaint.2018.03.019>

Finkel R.C., Owen L.A., Barnard P.L., Caffee M.W., 2003. *Beryllium-10 dating of Mount Everest moraines indicates a strong monsoon influence and glacial synchronicity throughout the Himalaya*. Geology, 31 (6), 561-564. [https://doi.org/10.1130/0091-7613\(2003\)031<0561:BDOMEM>2.0.CO;2](https://doi.org/10.1130/0091-7613(2003)031<0561:BDOMEM>2.0.CO;2)

Fu C. 1992. *Transitional climate zones and biome boundaries: a case study from China*. In: Hansen A.J., Castri, F.D. (Eds), *Landscape Boundaries*, 394-402. Springer Verlag, New York. [https://doi.org/10.1007/978-1-4612-2804-2\\_20](https://doi.org/10.1007/978-1-4612-2804-2_20)

Ganju A., Nagar Y.C., Sharma L.N., Sharma S., Juyal N., 2018. *Luminescence chronology and climatic implication of the late quaternary glaciation in the Nubra valley, Karakoram Himalaya, India*. Palaeogeography, Palaeoclimatology, Palaeoecology, 502, 52-62. <https://doi.org/10.1016/j.palaeo.2018.04.022>

Ganopolski A., Rahmstorf S., Petoukhov V., Claussen M., 1998. *Simulation of modern and glacial climates with a coupled global model of intermediate complexity*. Nature, 391 (6665), 351-356. <https://doi.org/10.1038/34839>

Gao Y., Cuo L., Zhang Y., 2014. *Changes in moisture flux over the Tibetan Plateau during 1979-2011 and possible mechanisms*. Journal of Climate, 27 (5), 1876-1893. <https://doi.org/10.1175/JCLI-D-13-00321.1>

Gayer E., Lavé J., Pik R., France-Lanord C., 2006. *Monsoonal forcing of Holocene glacier fluctuations in Ganesh Himal (Central Nepal) constrained by cosmogenic  $^{3}\text{He}$  exposure ages of garnets*. Earth and Planetary Science Letters, 252 (3-4), 275-288. <https://doi.org/10.1016/j.epsl.2006.09.040>

Ghosh R., Bera S., Sarkar A., Paruya D.K., Yao Y.F., Li C.S., 2015. *A ~50 ka record of monsoonal variability in the Darjeeling foothill region, eastern Himalayas*. Quaternary Science Reviews, 114, 100-115. <https://doi.org/10.1016/j.quascirev.2015.02.002>

Hahn D.G., Shukla J., 1976. *An apparent relationship between Eurasian snow cover and Indian monsoon rainfall*. Journal of the Atmospheric Sciences, 33 (12), 2461-2462. [https://doi.org/10.1175/1520-0469\(1976\)033<2461:AARBES>2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033<2461:AARBES>2.0.CO;2)

Heinrich H., 1988. *Origin and consequences of cyclic ice rafting in the northeast Atlantic Ocean during the past 130,000 years*. Quaternary research, 29 (2), 142-152. [https://doi:10.1016/0033-5894\(88\)90057-9](https://doi:10.1016/0033-5894(88)90057-9)

Herzschuh U., 2006. *Palaeo-moisture evolution in monsoonal Central Asia during the last 50,000 years*. Quaternary Science Reviews, 25 (1-2), 163-178. <https://doi.org/10.1016/j.quascirev.2005.02.006>

Hodell D.A., Nicholl J.A., Bontognali T.R., Danino S., Dorador J., Dowdeswell J.A., Einsle J., Kuhlmann H., Martrat B., Mleneck-Vautravers M.J., Rodríguez-Tovar F.J., 2017. *Anatomy of Heinrich Layer 1 and its role in the last deglaciation*. Paleoceanography, 32 (3), 284-303. <https://doi.org/10.1002/2016PA003028>

Hu G., Yi C.L., Zhang J.F., Liu J.H., Jiang T., 2015. *Luminescence dating of glacial deposits near the eastern Himalayan syntaxis using different grain-size fractions*. Quaternary Science Reviews, 124, 124-144. <https://doi.org/10.1016/j.quascirev.2015.07.018>

Hudson A.M., Quade J., 2013. *Long-term east-west asymmetry in monsoon rainfall on the Tibetan Plateau*. Geology, 41 (3), 351-354. <https://doi.org/10.1130/G33837.1>

Jia J., Fuyuan G., Dunsheng X., Wei H., Fahu C., 2018. *Moisture variations in arid central Asia and its out-of-phase relationship with the Asian monsoon during MIS 5: evidence from loess records*. Journal of Quaternary Science, 33 (4), 435-443. <https://doi.org/10.1002/jqs.3024>

Johnsen S.J., Clausen H.B., Dansgaard W., Gundestrup N.S., Hammer C.U., Andersen U., Andersen K.K., Hvidberg C.S., Dahl-Jensen D., Steffensen J.P., Shoji H., 1997. *The  $\delta^{18}\text{O}$  record along the Greenland Ice Core Project deep ice core and the problem of possible Eemian climatic instability*. Journal of Geophysical Research: Oceans, 102 (C12), 26397-26410. <https://doi.org/10.1029/97JC00167>

Kessler T.C., Klint K.E.S., Nilsson B., Bjerg P.L., 2012. *Characterization of sand lenses embedded in tills*. Quaternary Science Reviews, 53, 55-71. <https://doi.org/10.1016/j.quascirev.2012.08.011>

Krishna A.P., 2005. *Snow and glacier cover assessment in the high mountains of Sikkim Himalaya*. Hydrological Processes: An International Journal, 19 (12), 2375-2383. <https://doi.org/10.1002/hyp.5890>

Kudrass H.R., Hofmann A., Doose H., Emeis K., Erlenkeuser H., 2001. *Modulation and amplification of climatic changes in the Northern Hemisphere by the Indian summer monsoon during the past 80 ky*. Geology, 29 (1), 63-66. [https://doi.org/10.1130/0091-7613\(2001\)029<0063:MAAOCC>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0063:MAAOCC>2.0.CO;2)

ENVIS, 2016. Environmental Information System (ENVIS) Centre, Ministry of Environment and Forest, Govt. of India. Available at: [http://www.sikenvis.nic.in/database/naturalresources\\_790.aspx?format=print](http://www.sikenvis.nic.in/database/naturalresources_790.aspx?format=print).

Kumar P., Sharma M.C., Deswal S., Manna I., Chakraborty E., Prakash S., 2023. *Last Glacial Maximum and subsequent glacial chronology in the monsoon-dominated Sikkim Himalaya, India*. Palaeogeography, Palaeoclimatology, Palaeoecology, 617, 111480. <https://doi.org/10.1016/j.palaeo.2023.111480>

Lauterbach S., Andersen N., Wang Y.V., Blanz T., Larsen T., Schneider R.R., 2020. *An ~130 kyr record of surface water temperature and  $\delta^{18}\text{O}$  from the northern Bay of Bengal: investigating the linkage between Heinrich events and Weak Monsoon Intervals in Asia*. Paleoceanography and Paleoclimatology, 35 (2), e2019PA003646. <https://doi.org/10.1029/2019PA003646>

Li Y.X., Törnqvist T.E., Nevitt J.M., Kohl B., 2012. *Synchronizing a sea-level jump, final Lake Agassiz drainage, and abrupt cooling 8200 years ago*. Earth and Planetary Science Letters, 315, 41-50. <https://doi.org/10.1016/j.epsl.2011.05.034>

Li Y., Song Y., Lai Z., Han L., An Z., 2016. *Rapid and cyclic dust accumulation during MIS 2 in Central Asia inferred from loess OSL dating and grain-size analysis*. Scientific Reports, 6 (1), 1-6. <https://doi.org/10.1038/srep32365>

Liu S., Li J., Zhang H., Cao P., Mi B., Khokattiwong S., Kornkanitnan N., Shi X., 2020. Complex response of weathering intensity registered in the Andaman Sea sediments to the Indian Summer Monsoon over the last 40 kyr. *Marine Geology*, 426, 106206. <https://doi.org/10.1016/j.margeo.2020.106206>

Lutz A.F., Immerzeel W.W., Shrestha A.B., Bierkens M.F.P., 2014. Consistent increase in High Asia's runoff due to increasing glacier melt and precipitation. *Nature Climate Change*, 4 (7), 587-592. <https://doi.org/10.1038/nclimate2237>

Maher B.A., 2008. Holocene variability of the East Asian summer monsoon from Chinese cave records: a re-assessment. *The Holocene*, 18 (6), 861-866. <https://doi.org/10.1177/0959683608095569>

Maurer J.M., Schaefer J.M., Rupper S., Corley A., 2019. Acceleration of ice loss across the Himalayas over the past 40 years. *Science advances*, 5 (6), eaav7266. <https://doi.org/10.1126/sciadv.aav726>

Mehta M., Majeed Z., Dobhal D.P., Srivastava P., 2012. Geomorphological evidences of post-LGM glacial advancements in the Himalaya: a study from Chorabari Glacier, Garhwal Himalaya, India. *Journal of earth system science*, 121 (1), 149-163. <https://doi.org/10.1007/s12040-012-0155-0>

Milner A.M., Khamis K., Battin T.J., Brittain J.E., Barrand N.E., Füreder L., Cauvy-Fraunié S., Gislason G.M., Jacobsen D., Hannah D.M., Hodson A.J., 2017. Glacier shrinkage driving global changes in downstream systems. *Proceedings of the National Academy of Sciences*, 114 (37), 9770-9778. <https://doi.org/10.1073/pnas.1619807114>

Mortheast P., Ali S.N., 2014. Luminescence dating using quartz-for end-users. *Gondwana Geological Magazine*, 29, 1-10.

Mote P.W., Kaser G., 2007. The Shrinking Glaciers of Kilimanjaro: can Global Warming Be Blamed? The Kibo ice cap, a "poster child" of global climate change, is being starved of snowfall and depleted by solar radiation. *American Scientist*, 95 (4), 318-325.

Murari M.K., Owen L.A., Dortch, J.M., Caffee M.W., Dietsch C., Fuchs M., Haneberg W.C., Sharma M.C., Townsend-Small A., 2014. Timing and climatic drivers for glaciation across monsoon-influenced regions of the Himalayan-Tibetan orogen. *Quaternary Science Reviews*, 88, 159-182. <https://doi.org/10.1016/j.quascirev.2014.01.013>

Murray A.S., Wintle A.G. 2000. Luminescence dating of quartz using an improved single-aliquot regenerative-dose protocol. *Radiation measurements*, 32 (1), 57-73. [https://doi.org/10.1016/S1350-4487\(99\)00253-X](https://doi.org/10.1016/S1350-4487(99)00253-X)

Nagar Y.C., Ganju A., Satyawali P.K., Juyal N., 2013. Preliminary optical chronology suggests significant advance in Nubra valley glaciers during the Last Glacial Maximum. *Current Science*, 96-101.

Otto-Bliesner B.L., Brady E.C., Clauzet G., Tomas R., Levis S., Kothavala Z., 2006. Last glacial maximum and Holocene climate in CCSM3. *Journal of Climate*, 19 (11), 2526-2544. <https://doi.org/10.1175/JCLI3748.1>

Pang H., Hou S., Kaspari S., Mayewski P., Introne D., Masson-Delmotte V., Jouzel J., Li Z., He Y., Hong S., Qin D., 2012. Atmospheric circulation change in the central Himalayas indicated by a high-resolution ice core deuterium excess record. *Climate Research*, 53 (1), 1-12. <https://doi.org/10.3354/cr01090>

Peng X., Chen Y., Liu G., Liu B., Li Y., Liu Q., Han Y., Yang W., Cui Z., 2019. Late Quaternary glaciations in the Cogarbu valley, Bhutanese Himalaya. *Journal of Quaternary Science*, 34 (1), 40-50. <https://doi.org/10.1002/jqs.3079>

Phillips W.M., Sloan V.F., Shroder Jr J.F., Sharma P., Clarke M.L., Rennell H.M., 2000. Asynchronous glaciation at Nanga Parbat, northwestern Himalaya Mountains, Pakistan. *Geology*, 28 (5), 431-434. [https://doi.org/10.1130/0091-7613\(2000\)28<431:AGANPN>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<431:AGANPN>2.0.CO;2)

Prescott J.R., Hutton J.T., 1994. Cosmic ray contributions to dose rates for luminescence and ESR dating: large depths and long-term time variations. *Radiation measurements*, 23 (2-3), 497-500. [https://doi.org/10.1016/1350-4487\(94\)90086-8](https://doi.org/10.1016/1350-4487(94)90086-8)

Raina V.K., Srivastava D., 2008. *Glacier atlas of India*. Geological Survey of India Publications, 7 (1), GSI Bangalore.

Rana N., Sharma S., Ali S.N., Singh S., Shukla A.D., 2019. Investigating the sensitivity of glaciers to climate variability since the MIS-2 in the upper Ganga catchment (Saraswati valley), Central Himalaya. *Geomorphology*, 346, 106854. <https://doi.org/10.1016/j.geomorph.2019.106854>

Rashid H., England E., Thompson L., Polyak L., 2011. Late glacial to Holocene Indian summer monsoon variability based upon sediment records taken from the Bay of Bengal. *TAO: Terrestrial, Atmospheric and Oceanic Sciences*, 22 (2), 2. <https://doi.org/10.3319/TAO.2010.09.17.02>

Rawat S., Gupta A.K., Sangode S.J., Srivastava P., Nainwal H.C., 2015. Late Pleistocene-Holocene vegetation and Indian summer monsoon record from the Lahaul, northwest Himalaya, India. *Quaternary Science Reviews*, 114, 167-181. <https://doi.org/10.1016/j.quascirev.2015.01.032>

Rohling E.J., Pälike H., 2005. Centennial-scale climate cooling with a sudden cold event around 8,200 years ago. *Nature*, 434 (7036), 975-979. <https://doi.org/10.1038/nature03421>

Rupper S., Roe G., Gillespie A., 2009. Spatial patterns of Holocene glacier advance and retreat in Central Asia. *Quaternary Research*, 72 (3), 337-346. <https://doi.org/10.1016/j.yqres.2009.03.007>

Sati S.P., Ali S.N., Rana N., Bhattacharya F., Bhushan R., Shukla A.D., Sundriyal Y., Juyal N., 2014. Timing and extent of Holocene glaciations in the monsoon-dominated Dunagiri valley (Bangni glacier), Central Himalaya, India. *Journal of Asian Earth Sciences*, 91, 125-136. <https://doi.org/10.1016/j.jseas.2014.05.008>

Scherler D., Bookhagen B., Strecker M.R., von Blanckenburg F., Rood D., 2010. Timing and extent of late Quaternary glaciation in the western Himalaya constrained by  $^{10}\text{Be}$  moraine dating in Garhwal, India. *Quaternary Science Reviews*, 29 (7-8), 815-831. <https://doi.org/10.1016/j.quascirev.2009.11.031>

Sharma S., Shukla A.D., 2018. Factors governing the pattern of glacier advances since the Last Glacial Maxima in the transitional climate zone of the Southern Zanskar Ranges, NW Himalaya. *Quaternary Science Reviews*, 201, 223-240. <https://doi.org/10.1016/j.quascirev.2018.10.006>

Shi Y.F., Kong Z.C., Wang S.M., Tang L.Y., Wang F.B., Yao T.D., Zhao X., Zhang P.Y., Shi S., 1994. Climates and environments of the Holocene mega-thermal maximum in China. *Science in China (Series B)* 37, 481-493. <https://doi.org/10.1360/yb1994-37-4-481>

Shukla T., Mehta M., Jaiswal M.K., Srivastava P., Dobhal D.P., Nainwal H.C., Singh A.K., 2018. Late Quaternary glaciation history of monsoon-dominated Dingad basin, central Himalaya, India. *Quaternary Science Reviews*, 181, 43-64. <https://doi.org/10.1016/j.quascirev.2017.11.032>

Sinha A., Cannariato K.G., Stott L.D., Li H.C., You C.F., Cheng H., Edwards R.L., Singh I.B., 2005. Variability of Southwest Indian summer monsoon precipitation during the Bølling-Allerød. *Geology*, 33 (10), 813-816. <https://doi.org/10.1130/G21498.1>

Sirocko F., Sarnthein M., Erlenkeuser H., Lange H., Arnold M., Dulessy J.C., 1993. Century-scale events in monsoonal climate over the past 24,000 years. *Nature*, 364 (6435), 322-324. <https://doi.org/10.1038/364322a0>

Soheb M., Ramanathan A.L., Pandey P., Mandal A., 2015. Climate change from Himalayan Glaciers' perspective. Case studies from India. *Journal of Climate Change*, 1 (1-2), 27-35. <https://doi.org/10.3233/JCC-150002>

Solomina O., Haeberli W., Kull C., Wiles G., 2008. *Historical and Holocene glacier-climate variations: general concepts and overview*. Global and Planetary Change, 60 (1), 1-9. <https://doi:10.1016/j.gloplacha.2007.02.001>

Steinhilber F., Beer J., Fröhlich C., 2009. *Total solar irradiance during the Holocene*. Geophysical Research Letters, 36 (19). <https://doi.org/10.1029/2009GL040142>

Thompson L.O., Yao T., Davis M.E., Henderson K.A., Mosley-Thompson E., Lin P.N., Bolzan J.F., 1997. *Tropical climate instability: The last glacial cycle from a Qinghai-Tibetan ice core*. Science, 276 (5320), 1821-1825. <https://doi.org/10.1126/science.276.5320.1821>

Tian L., Yao T., MacClune K., White J.W.C., Schilla A., Vaughn B., Vachon R., Ichiyanagi K., 2007. *Stable isotopic variations in west China: a consideration of moisture sources*. Journal of Geophysical Research: Atmospheres, 112 (D10). <https://doi.org/10.1029/2006JD007718>

Tierney J.E., Pausata F.S., Demenocal P., 2016. *Deglacial Indian monsoon failure and North Atlantic stadials linked by Indian Ocean surface cooling*. Nature Geoscience, 9 (1), 46-50. <https://doi.org/10.1038/ngeo2603>

Tsukamoto S., Asahi K., Watanabe T., Rink W.J., 2002. *Timing of past glaciations in Kanchenjunga Himalaya, Nepal by optically stimulated luminescence dating of tills*. Quaternary International, 97, 57-67. [https://doi.org/10.1016/S1040-6182\(02\)00051-4](https://doi.org/10.1016/S1040-6182(02)00051-4)

Wang Q., Yi S., Chang L., Sun W., 2017. *Large-scale seasonal changes in glacier thickness across High Mountain Asia*. Geophysical Research Letters, 44 (20), 10-427. <https://doi.org/10.1002/2017GL075300>

Wang Y.J., Cheng H., Edwards R.L., An Z.S., Wu J.Y., Shen C.C., Dorale J.A., 2001. *A high-resolution absolute-dated late Pleistocene monsoon record from Hulu Cave, China*. Science, 294 (5550), 2345-2348. <https://doi.org/10.1126/science.1064618>

Wang Y., Cheng H., Edwards R.L., He Y., Kong X., An Z., Wu J., Kelly M.J., Dykoski C.A., Li X., 2005. *The Holocene Asian monsoon: links to solar changes and North Atlantic climate*. Science, 308 (5723), 854-857. <https://doi.org/10.1126/science.1106296>

Wei K., Gasse F., 1999. *Oxygen isotopes in lacustrine carbonates of West China revisited: implications for post glacial changes in summer monsoon circulation*. Quaternary Science Reviews, 18 (12), 1315-1334. [https://doi.org/10.1016/S0277-3791\(98\)00115-2](https://doi.org/10.1016/S0277-3791(98)00115-2)

Yan Q., Owen L.A., Wang H., Zhang Z., 2018. *Climate constraints on glaciation over High-Mountain Asia during the last glacial maximum*. Geophysical Research Letters, 45 (17), 9024-9033. <https://doi.org/10.1029/2018GL079168>

Yang S., Ding Z., 2014. *A 249 kyr stack of eight loess grain size records from northern China documenting millennial-scale climate variability*. Geochemistry, Geophysics, Geosystems, 15 (3), 798-814. <https://doi.org/10.1002/2013GC005113>

Yang W., Chu W., Liu G., 2022. *Importance of the seasonal temperature and precipitation variation on glacial evolutions during the LGM at monsoonal Himalaya based on Cogarbu valley*. Palaeogeography, Palaeoclimatology, Palaeoecology, 601, 111132. <https://doi.org/10.1016/j.palaeo.2022.111132>

Yihui D., Zunya W., 2008. *A study of rainy seasons in China*. Meteorology and Atmospheric Physics, 100 (1), 121-138. <https://doi.org/10.1007/s00703-008-0299-2>

Yuan D., Cheng H., Edwards R.L., Dykoski C.A., Kelly M.J., Zhang M., Qing J., Lin Y., Wang Y., Wu J., Dorale J.A., 2004. *Timing, duration, and transitions of the last interglacial Asian monsoon*. Science, 304 (5670), 575-578. <https://doi.org/10.1126/science.1091220>

Zech R., Zech M., Kubik P.W., Kharki K., Zech W., 2009. *Deglaciation and landscape history around Annapurna, Nepal, based on 10Be surface exposure dating*. Quaternary Science Reviews, 28 (11-12), 1106-1118. <https://doi.org/10.1016/j.quascirev.2008.11.013>

Zhou L., Chen H., Hua W., Dai Y., Wei N., 2016. *Mechanisms for stronger warming over drier ecoregions observed since 1979*. Climate Dynamics, 47, 2955-2974. <https://doi.org/10.1007/s00382-016-3007-9>

(Ms. received 19 July 2024, accepted 06 March 2025)

